arXiv:1704.01431v1 [astro-ph.SR] 5 Apr 2017

Late-time flattening of Type Ia Supernova light curves: Constraints from SN 2014J in M82

Yi Yang[1], Lifan Wang[1,2], Dietrich Baade[3], Peter. J. Brown[1], Misty Cracraft[4], Peter A. Höflich[5], Justyn Maund[6,†], Ferdinando Patat[3], William B. Sparks[4], Jason Spyromilio[3], Heloise F. Stevance[6], Xiaofeng Wang[7], J. Craig Wheeler[8]

Received _____; accepted _____

[1]George P. and Cynthia Woods Mitchell Institute for Fundamental Physics & Astronomy, Texas A. & M. University, Department of Physics and Astronomy, 4242 TAMU, College Station, TX 77843, USA, email: ngc4594@physics.tamu.edu

[2]Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210008, China

[3]European Organisation for Astronomical Research in the Southern Hemisphere (ESO), Karl-Schwarzschild-Str. 2, 85748 Garching b. München, Germany

[4]Space Telescope Science Institute, Baltimore, MD 21218, USA

[5]Department of Physics, Florida State University, Tallahassee, Florida 32306-4350, USA

[6]Department of Physics and Astronomy, University of Sheffield, Hicks Building, Hounsfield Road, Sheffield S3 7RH, UK

[7]Physics Department and Tsinghua Center for Astrophysics (THCA), Tsinghua University, Beijing, 100084, China

[8]Department of Astronomy and McDonald Observatory, The University of Texas at Austin, Austin, TX 78712, USA

[†]Royal Society Research Fellow

ABSTRACT

The very nearby Type Ia supernova 2014J in M82 offers a rare opportunity to study the physics of thermonuclear supernovae at extremely late phases ($\gtrsim$800 days). Using the *Hubble Space Telescope*, we obtained five epochs of high precision photometry for SN 2014J from 277 days to 985 days past the $B-$band maximum light. The reprocessing of electrons and X-rays emitted by the radioactive decay chain $^{57}\mathrm{Co}\rightarrow^{57}\mathrm{Fe}$ are needed to explain the significant flattening of both the $F606W$-band and the pseudo-bolometric light curves. The flattening confirms previous predictions that the late-time evolution of type Ia supernova luminosities requires additional energy input from the decay of ^{57}Co (Seitenzahl et al. 2009). By assuming the $F606W$-band luminosity scales with the bolometric luminosity at $\sim$500 days after the $B-$band maximum light, a mass ratio $^{57}\mathrm{Ni}/^{56}\mathrm{Ni}\sim0.076^{+0.009}_{-0.008}$ is required. This mass ratio is roughly $\sim$3.5 times the solar ratio and favors a progenitor white dwarf with a mass near the Chandrasekhar limit. A similar fit using the constructed pseudo-bolometric luminosity gives a mass ratio $^{57}\mathrm{Ni}/^{56}\mathrm{Ni}\sim0.094^{+0.017}_{-0.016}$. Astrometric tests based on the multi-epoch *HST* ACS/WFC images reveal no significant circumstellar light echoes in between 0.3 pc and 100 pc (Yang et al. 2017) from the supernova 2014J.

Subject headings: abundances — nuclear reactions — nucleosynthesis — supernovae: individual (SN 2014J)

1. Introduction

The astronomical community widely agrees that luminous hydrogen-poor Type Ia supernovae (SNe) explosions are powered by the thermonuclear runaway of ($\geqslant 1M_{\odot}$) carbon/oxygen white dwarfs (WDs Hoyle & Fowler 1960). The accretion-induced explosion fuses $\sim$0.1-1.0$M_{\odot}$ of radioactive ^{56}Ni. Type Ia SNe cosmology, which uses Type Ia SNe as the most accurate distance indicators at redshifts out to $z \sim$2 (Riess et al. 1998; Perlmutter et al. 1999; Riess et al. 2016). Amazingly, this accuracy is achieved without knowing exactly the nature of various progenitors.

Prior to maximum luminosity, the light curve of Type Ia SNe is powered by the energy generated by the decay of explosion-synthesized radioactive nuclei. The reprocessing in the ejecta converts the energy to longer wavelengths. The decay chain of ^{56}Ni$\rightarrow^{56}$Co$\rightarrow^{56}$Fe provides the main source of energy deposition into the ejecta of Type I SNe (Arnett 1982). During the early phases, the optically-thick ejecta trap the energy. The dominant process is Compton scattering of γ-rays produced by the decay $^{56}\mathrm{Ni} + e^{-} \rightarrow ^{56}\mathrm{Co} + \gamma + \nu_e$, ($t_{1/2} \sim$6.08 days), which allows energy to escape as X-ray continuum or absorbed by the material in the ejecta via the photoelectric effect (see Milne et al. 1999; Penney & Hoeflich 2014 for comprehensive reviews). The produced ^{56}Co decays to stable ^{56}Fe, and the ^{56}Co decay process, with half-life $t_{1/2} \sim$77 days, dominates after $\sim$200 days, when the expanding ejecta become more and more optically thin, and the column density decreases as t^{-2} (e.g., Arnett 1979; Chan & Lingenfelter 1993; Cappellaro et al. 1997; Milne et al. 1999). The ^{56}Co decays Eighty-one percent of the ^{56}Co decays as follows: electron capture: $^{56}\mathrm{Co} + e^{-} \rightarrow ^{56}\mathrm{Fe} + \gamma + \nu_e$, and the remaining through annihilation of high energy positrons in the ejecta, $^{56}\mathrm{Co} \rightarrow ^{56}\mathrm{Fe} + e^{+} + \gamma + \nu_e$.

Observations at extremely late phases provide unique opportunities to examine various models exploring the effects of a magnetic field. As long as energy deposition is dominated

by positrons being completely trapped by the magnetic field, the slope of the bolometric light curve should match the ^{56}Co decay rate. On the other hand, Milne et al. (1999) suggested a "radially combed" magnetic field, or even a magnetic-field-free situation as no magnetic field in radial directions will leads to an increasing fraction of positron escape, and the light curve should decline faster than the rate of ^{56}Co decay. The discrepancy between the "trapping scenario" with a confining magnetic field and the case without magnetic field can be as significant as 2 magnitudes in the photometric light curves from 400 - 800 days (see Figure 9 of Milne et al. 1999). Similar variations of the late-time light curves have been found by Penney & Hoeflich (2014) based on measuring positron transport effects and their dependency on the magnetic field with late-time line profiles. However, they drew different conclusions. As the SN envelope undergoes homologous expansion, the morphology of the magnetic field remains, but the Lamor radius increased linearly with time. Therefore, the fraction of escaped photons would exhibit a time-dependence due to the variations of the magnetic field. The light curve should decline faster than the rate of ^{56}Co decay.

Additonally, different effects of nucleosynthesis can be testable through the very late photometric evolution of Type Ia SNe and may be used to discriminate between different explosion models. Two of the most favorable explosion channels: a delayed detonation in a Chandrasekhar-mass white dwarf (Khokhlov 1991) and a violent merger of two carbon-oxygen white dwarfs (Pakmor et al. 2011, 2012), will result in late-time light curves behaving differently due to different amounts of ejecta heating from ^{57}Co and ^{55}Fe (Röpke et al. 2012). Therefore, fitting the decline rate of the light curve at extremely late times provides a unique opportunity to test the enigmatic explosion mechanisms of Type Ia SNe.

Increasing evidence shows the flattening of Type Ia SN light curves around 800 to 1000 days, i.e., SN 1992A ($\sim$950 days; Cappellaro et al. 1997, Cappellaro et al. 1997), SN 2003hv ($\sim$700 days; Leloudas et al. 2009), and SN 2011fe ($\sim$930 days; Kerzendorf et al. 2014). This

flattening cannot be explained even by complete trapping of the ^{56}Co positrons. Seitenzahl et al. (2009) suggested that additional heating from the Auger and internal conversion electrons, together with the associated X-ray cascade produced by the decay of ^{57}Co$\rightarrow^{57}$Fe ($t_{1/2}$ ≈272 days) and ^{55}Fe$\rightarrow^{55}$Mn ($t_{1/2}$ ≈1000 days), will significantly slow down the decline of the light curve.

Only very recently, Graur et al. (2016) carried out an analysis of the light curve of SN 2012cg as late as $\sim$ 1055 days after the explosion and excluded the scenario in which the light curve of SN 2012cg is solely powered by the radioactive decay chain ^{56}Ni$\rightarrow^{56}$Co$\rightarrow^{56}$Fe, unless there is an unresolved light echo $\sim$14 magnitudes fainter than the SN peak luminosity. Another very careful study on the late-time evolution of SN 2011fe has already extended the observing effort to an unprecedented 1622 days past the $B-$band maximum light (Shappee et al. 2016). This analysis has clearly detected the radioactive decay channel powered by ^{57}Co, with a mass ratio of $\log(^{57}\mathrm{Co}/^{56}\mathrm{Co}) = -1.62^{+0.08}_{-0.09}$. This abundance ratio is strongly favored by double degenerate models which require a lower central density. The detection of ^{55}Fe is still unclear at these late epochs (Shappee et al. 2016). Another study based on the pseudo-bolometric light curve for the SN 2011fe has measured the mass ratio of ^{57}Co to ^{56}Co to be 1.3 – 2.5 times the solar value, which is broadly consistent with the ratios predicted for the delayed detonation models (Dimitriadis et al. 2017). Additionally, spectroscopic information of the nearby SN 2011fe has been obtained at 981 days (Graham et al. 2015) and 1034 days (Taubenberger et al. 2015). Strong energy input from the radioactive decay of ^{57}Co is required, without which the optical spectrum would be underproduced by a factor of $\sim$4 (Fransson & Jerkstrand 2015). The mass ratio of ^{57}Ni to ^{56}Ni produced, which gives a strong constraint on the Type Ia SN explosions, is found to be roughly 2.8 and 2 times of the solar ratio for SN 2011fe and SN 2012cg, respectively (Fransson & Jerkstrand 2015; Graur et al. 2016).

SN 2014J was first discovered on Jan 21.805 UT by Fossey et al. (2014) in the very nearby starburst galaxy M82 (3.53±0.04 Mpc, Dalcanton et al. 2009). Later observations constrained the first light of the SN to Jan. 14.75 UT (Zheng et al. 2014; Goobar et al. 2014). This date is consistent with the early rising recorded by the 0.5-m Antarctic Survey Telescope (AST) during its test observations (Ma et al. 2014) as well as with other pre-discovery limits reported by various groups (Denisenko et al. 2014; Itagaki et al. 2014; Gerke et al. 2014). SN 2014J reached its $B-$band maximum on Feb. 2.0 UT (JD 2,456,690.5) at a magnitude of 11.85±0.02 (Foley et al. 2014). Follow-up photometric and spectroscopic observations have been made by various groups (Johansson et al. 2014; Lundqvist et al. 2015; Bonanos & Boumis 2016; Srivastav et al. 2016). The strength of γ-ray lines (Churazov et al. 2014; Diehl et al. 2015) and an analytic model fit to the pseudo bolometric light curve (Srivastav et al. 2016) of SN 2014J suggest that ∼0.5-0.6 $M_{\odot}$ of ^{56}Ni has been synthesized in the explosion. In this paper, we present our late time *HST* photometric observations of SN 2014J and fit both the $F606W$ (broad V) band and an estimate of the pseudo-bolometric luminosity evolution with the Bateman equation considering the luminosity contributed by the decay of ^{56}Co, ^{57}Co, and ^{55}Fe. In addition to following a similar approach presented in Graur et al. (2016), we provide a careful astrometric analysis to the time-evolution of the position and profile of the SN 2014J point source at very late epochs.

2. Observations and Data Reduction

We imaged the SN 2014J with the *Hubble Space Telescope* Advanced Camera for Surveys/Wide Field Channel (*HST* ACS/WFC) during five visits (V1-V5) under multiple *HST* programs: GO-13717 (PI: Wang), GO-14139 (PI: Wang), and GO-14663 (PI: Wang), i.e., V1∼day 277, V2∼day 416, V3∼day 649, V4∼ day 796, and V5∼ day 983, relative to

its $B-$band maximum at a mgnitude of 11.85±0.02 on Feb. 2.0 UT (JD 2,456,690.5, Foley et al. 2014). Figure 1 shows the field around SN 2014J. A log of observations is presented in Table 1. Exposures obtained with different ACS visual polarizers and in different filter combinations and visits have been aligned through *Tweakreg* in the *Astrodrizzle* package (Gonzaga et al. 2012).

The throughput of each ACS/WFC polarizer being used by the *Synphot* [1] software does not match the values determined by on-orbit calibrations. We corrected the polarizers' throughput with the values deduced by on-orbit calibrations (i.e., Table 12 of Cracraft & Sparks 2007, also see Biretta et al. 2004). Following the three polarizers case described in earlier works by Sparks & Axon (1999), we deduced the Stokes vectors from the observations. In this work, we only discuss the observed flux from the SN 2014J, and the intensity maps (Stokes I) are the only required input parameter to this analysis.

$$I = \frac{2}{3}[r(POL0) + r(POL60) + r(POL120)], \tag{1}$$

where $r(POL0)$, etc. are the count rates in the images obtained through the three polarizers. The polarimetric properties of the SN 2014J among different late phases will be discussed in a future work.

After ∼600 days past maximum light, the SN became sufficiently dim and the count rates at the central pixels of the SN PSF became comparable to the bright part of the nebulosity close to the SN. The field shows that the SN lies at one end of a dark lane, and just west of a bright patch of nebulosity. A background subtraction procedure significantly diminishes the time-invariant signals and improves the photometry of evolving faint sources. Unfortunately, we found no pre-SN Hubble images, either with or without the polarizers, showing the same region using filters compatible with our observations. Images obtained

[1]http://www.stsci.edu/institute/software_hardware/stsdas/synphot

on March 29 2006 (program #10776; PI:Mountain) with *HST* ACS/WFC in the $F435W$, $F555W$, and $F814W$ were used as background templates for our $F475W$, $F606W$, and $F775W$ exposures, respectively. For each band, the background templates have been scaled and subtracted from the intensity map. Measurements were made with a circular aperture of 0.1″ (2 pixels in the ACS/WFC FOV) with aperture corrections according to Hartig (2009) and Sirianni et al. (2005). The photometry has been performed using the IRAF [2] APPHOT package. The residual of the background has been estimated by the median pixel value of an annulus around the SN. Compromising between determining the local background residual with nearby pixels and excluding the contamination from resolved interstellar light echoes (Yang et al. 2017), we choose the inner and outer radii as 1.2″ (24 pixels) and 1.5″ (30 pixels) for V1 and V2, and 0.6″ (12 pixels) and 0.8″ (16 pixels) for V3, V4, and V5. Table 2 presents the AB magnitude of SN 2014J at five late epochs. The photometric uncertainties include the Poisson noise of the signal, the photon noise of the background, the readout noise contribution (3.75 electrons/pixel for ACS/WFC), and the uncertainties in aperture corrections. These quantities were added in quadrature.

This photometry strategy has been carried out considering that extremely nonuniform background structures dominate the error budget in the late phases of the SN 2014J photometry, especially in V4 and V5. For the scientific consideration of this study, which is testing the models for the light curve evolution at very late phases, the major concern in the data reduction procedure is to obtain the correct decline rate of the SN light curves. Therefore, we conducted a sanity check to test the reliability of our measurement by performing photometry on differenced images from our observations obtained at different

[2]IRAF is distributed by the National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation (NSF).

epochs. This directly measures the differential fluxes and therefore the light curve decline rate. The decline rates between all the epochs calculated from photometry shown in Table 3 and measured by this sanity check agree within ∼2% and are smaller than the photometric uncertainties.

We correct our measurements for both the interstellar dust extinction in the SN host galaxy and the Galactic extinction towards the SN 2014J. In fact, any imperfection in the extinction correction will only affect the individual magnitudes but no the decaline rates of the light curves. A peculiar extinction law R_V ∼1.4 towards the SN 2014J line of sight has been suggested by many studies (Amanullah et al. 2014; Brown et al. 2015; Foley et al. 2014; Gao et al. 2015; Goobar et al. 2014). In this study, we adopt $R_V = 1.44\pm0.03$ and $A_V = 2.07\pm0.18$ from Foley et al. (2014) for the extinction from the host galaxy and $R_V = 3.1$ and $E(B-V) = 0.054$ mag for the Galactic extinction following Foley et al. (2014) based on Dalcanton et al. (2009) and Schlafly & Finkbeiner (2011). Extinction in $F475W$, $F606W$, and $F775W$ has been calculated for each component using a reddening law from Cardelli et al. (1989) with the corresponding R_V value. Both components are added to account for the total extinction towards SN 2014J for each *HST* ACS bandpass.

3. Analysis

In this section, we will test different mechanisms powering the late-time light curve, and whether the light curve behavior is consistent with the prediction for the delayed-detonation and the violent merger progenitor scenarios following a similar procedure as Graur et al. (2016) for SN 2012cg. We assume that the ejecta do not interact with any circumstellar material. The pseudo-bolometric light curve for SN 2014J was calculated over a wavelength range from 3500Å - 9000Å based on our multi-band optical photometry. We briefly summarize the steps as follows:

(1) Based on the lack of significant spectral evolution of SN 2011fe compared to a spectrum at 593 days (Graham et al. 2015), we assume the MODS/LBT spectrum of SN 2011fe at 1016 days (Taubenberger et al. 2015) represents the major spectral features of SN 2014J on V3∼day 649, V4∼day 796, and V5∼day 983. The spectrum was downloaded from the WISeREP archive [3].

(2) We then perform synthetic photometry from this spectrum for the $F475W$, $F606W$, and $F775W$ bands.

(3) We calculate the differences between the synthetic photometry of the SN 2011fe spectrum and our extinction-corrected, observed photometry of SN 2014J.

(4) We calculate the scale factors between the observed and synthetic magnitudes in each filter.

(5a) We warp the spectrum using a $2^{\rm nd}$ order polynomial fit to the scale factors at the effective wavelength [4].

(5b) Alternatively, for each epoch, we fit a single wavelength-independent gray scale across all wavelengths.

(6) We iterate steps (2) - (5) until the synthetic and observed photometry match to better than 0.02 mag in each filter for (5a), or the mean difference between the synthetic and the observed photometry converges to its minimum value for (5b), which standard deviation among the three filters gives 0.11 mag.

The pseudo-bolometric luminosity for each epoch was obtained by integrating the scaled spectrum returned from (5a) or (5b) over the wavelength range from 3500Å - 9000Å. The errors on the pseudo-bolometric light curve were computed through a Monte Carlo re-sampling approach using photometric errors. The warping in (5a) aims at iteratively

[3]http://wiserep.weizmann.ac.il

[4]http://pysynphot.readthedocs.io/en/latest/properties.html#pysynphot-formula-efflam

producing spectra consistent with the photometry which follows a very similar procedure as described in Shappee et al. (2016), while the scaling in (5b) is less sensitive to the extrapolation of the polynomial correction to the spectrum. The pseudo-bolometric luminosities calculated from these two approaches agree within 6% at all epochs, compatible with the uncertainties of the Monte Carlo approach. Our analysis of the bolometric evolution of SN 2014J is based on the bolometric luminosity obtained with (5b). Qualitatively similar results have been obtained by duplicating the entire analysis based on (5a) as follows.

In Figure 2 we present the spectra with the warping procedure (left panel) and with the gray scaling (right panel). For comparison, in each upper panel, we overplotted the bandpass monochromatic flux calculated as the product Total Counts × PHOTFLAM [5], where PHOTFLAM is the inverse sensitivity (in erg cm^{-2} s^{-1} Å^{-1}) representing a signal of 1 electron per second. The lower panels present the total bandpass throughput curve (*HST* + ACS) [6] for our $F475W$, $F606W$, and $F775W$ observations. The spectra on the left panel are iterated to agree quantitatively with the photometry. Visual differences between the monochromatic bandpass flux and the spectra arise because PHOTFLAM used for the SED assumes a smooth AB spectrum, which is different than the SN spectrum (see Brown et al. 2016 for a comprehensive discussion).

In the left panels of Figure 3, we present the $F475W$, $F606W$, and $F775W$-band luminosity of SN 2014J after the extinction correction. In addition to fitting the pseudo-bolometric light curve after ∼500 days with the contribution from three decay chains: ^{56}Co→^{56}Fe, ^{57}Co→^{57}Fe, and ^{55}Fe→^{55}Mn (an 'all isotopes' model), we also fit the same model to our $F606W$-band observations. Here we have assumed that after ∼500 days

[5]This can be obtained with the ACS Zeropoints Calculator at https://acszeropoints.stsci.edu/

[6]http://www.stsci.edu/hst/acs/analysis/throughputs

the $F606W$-band, which is centered at wavelength 5888.8Å and with a width [7] of 2570$\mathring{A}$, captures the dominant Fe features ([Fe II] around 4700Å and 5300Å, blended [Fe II]λ7155 and [Ni II]λ7378 around 7200Å; Taubenberger et al. 2015) and to be proportional to the bolometric light curves as V-band observations (Milne et al. 2001).

Limited by a small number of visits, we approximate the 'all isotopes' model with two free parameters: the mass ratio $M(^{57}\mathrm{Co})/M(^{56}\mathrm{Co})$, and a scale factor to match the $F606W$ photometry (or the pseudo-bolometric luminosity) with the model-calculated values. Using the solution to the Bateman equation which describes the abundances and activities in a decay chain as a function of time (following Seitenzahl et al. 2014), and by counting the decay energy carried by charged leptons and X-rays, the luminosity contribution from a single decay chain gives:

$$L_A(t) = 2.221 \frac{C}{A} \frac{\lambda_A}{\mathrm{days}^{-1}} \frac{M(A)}{M_\odot} \frac{q_A^l + q_A^X}{\mathrm{keV}} \exp(-\lambda_A t_e) \times 10^{43} \mathrm{ergs}^{-1} \quad (2)$$

where C is a scaling factor, A gives the corresponding atomic number, λ_A is the inverse mean lifetime which is $\lambda_A = \tau_A^{-1} = \ln(2)/t_{1/2,A}$, $M(A)$ is the total mass of a certain decaying element, q_A^l and q_A^X are the average energies per decay carried by charged leptons and X-rays, respectively, and t_e is the time since explosion. Due to the limited data points in our late-time photometry, we used a ratio of $M(^{57}Co)/M(^{55}Fe) \approx 0.8$ (model rpc32; Ohlmann et al. 2014). The values of λ_A, q_A^l and q_A^X used here are obtained from Table 1 of Seitenzahl et al. (2009) and Table 2 of Seitenzahl et al. (2014). We justify our assumptions as follows: (1) The total deposition function is determined by both the net deposition functions for $\gamma-$rays and positrons. The $\gamma-$rays produced by the annihilation of the positrons are subject to both deposition functions. By simply assuming the radioactive source is confined to the center of a spherical distribution of ejecta yields a fraction $1 - e^{-\tau_\gamma}$ of the energy produced by $\gamma-$rays would be left behind in the ejecta (Swartz & Wheeler 1991). The

[7] where the filter throughput is larger than 0.05%

$\gamma-$ray optical depth τ_γ drops significantly as t^{-2} and we neglect contributions from γ-rays because the SN ejecta became transparent to γ-rays at $t \gtrsim$500 days (Milne et al. 2001); (2) Limited by a small number of photometric point, we begin by fitting Equation 2 assuming full trapping of positrons/electrons. In other words, we assume positrons, electrons, and X-rays are fully trapped, instantaneously deposited, and radiate their energy. One should also note that very recently, Dimitriadis et al. (2017) found that the late-time bolometric light curve of SN 2011fe is consistent with both models, namely positron/electron escape as well as complete trapping, that allow for positron/electron escape, and models that have complete positron/electron trapping but do allow for redistribution of flux to the mid-far IR.

The luminosity contribution from each decay channel is shown in Figure 3. The total luminosity given by these decay chains is represented by the pink dashed line. In the left panel, we show that a mass ratio of $M(^{57}\mathrm{Co})/M(^{56}\mathrm{Co}) = 0.076^{+0.009}_{-0.008}$ gives the best fit to the 'all isotopes' model based on the $F606W$-band observations after $t \sim 500$ days (V3, V4, and V5). The dot-dashed gray lines show the model including the luminosity from ^{56}Co decay and possible reflections from an unresolved t^{-1} light echo (see Graur et al. 2016). In the right panel, we show the same trend in a similar fitting based on the pseudo-bolometric light curve, which the mass ratio gives $M(^{57}\mathrm{Co})/M(^{56}\mathrm{Co}) = 0.094^{+0.017}_{-0.016}$.

If light echoes dominate the late time signal from the SN, we may expect a significant profile change or centroid drift if the circumstellar matter is distributed at sufficiently large distances from the SN. Light scattered by dust at such distances can produce measurable distortions to the image profiles if the scattered light dominates the total observed flux. At the distance of SN 2014J, 1 light year corresponds to 0.17 *HST* ACS/WFC pixels. Depending on the dust distribution, we may expect the stellar profiles to become non-point like, or the centroid of the stellar profile to drift at late time. We have checked the

stellar profiles and found no significant deviations from a point source at all epochs of our observations. In the following, we provide a comprehensive check on the centroid position of the SN.

The barycenter of the stars and HII regions around SN 2014J were measured to estimate a possible change in the relative position of the light emission of the SN. The precision is limited by the scarcity of stars in the immediate vicinity of the SN, as well as the uncharacterized field distortions caused by ACS/WFC polarizers (see, i.e., Section 5.3 of Gonzaga et al. 2012). Figure 4 presents the apparent shift in position measured from our observations in $F475W$ and $F606W$. The RA and Dec were calculated using the image from V3, with the SN at the origin of the coordinates. The gray arrows show the vector difference of the originally measured positions of the star on two different epochs. The black arrow shows the same vector after a 2-D linear regression to remove the dependence on RA and Dec, which may be caused by residual errors of astrometric calibrations. The linear regression was found to be able to reduce the shift significantly in all cases. The reference objects for astrometric comparisons were selected within a radius of 500 pixels of the position of the SN. The FWHM of the objects was restricted to be less than 8 pixels. Only a small number objects in the earliest V1 satisfy these criteria due to the relatively short exposure time.

In Figure 4, the upper panels present the measurements based on the highest S/N $F475W$-band exposures, and the lower panels present the same figures for $F606W$. For instance, the upper-left panel presents the comparison between V3 and V1. The SN exhibits an apparent motion of $0.050''$; after linear regression with the RA and Dec, this reduces to $0.021''$. This is in agreement with all the other objects in the field, which show an average distance shift of $0.015''$ and an RMS of $0.013''$. The upper-middle panel presents the comparison between V3 and V2. The SN exhibits an apparent drift in position of $0.019''$;

after linear regression this reduces to 0.005″. The field objects exhibit an average drift of 0.013″ and an RMS of 0.009″. This also implies that there is no apparent position drift of the SN with an upper limit of 0.014″. The upper-right panel presents the comparison between V3 and V4. The original images exhibit a position shift of 0.009″ for the SN. This is already too small to be a significant drift. After linear regression with RA and Dec, using the stars around the SN, the drift changes to 0.010″. The mean drift of objects in the field is 0.012″ with an RMS of 0.007″. An upper bound on the centroid position drift of the SN between V3 and V4 is thus observed to be 0.017″. Similar results were obtained for $F606W$-band exposures. In all cases, we have not observed a significant position drift of the SN. The absence of such drift sets a strong constraint on the nature of the late time emission from SN 2014J. If the significant flattening in $F606W$ is due to light echoes, the dust must be lie within 0.017″ of the SN.

4. Discussion and Summary

Table 3 shows the decline rate of the light curves at different epochs. Before t ∼600 days, the SN dims more rapidly than the light curve powered solely by the ^{56}Co decay. The γ-ray energy deposition becomes no longer significant after ∼200 days, therefore, a substantial fraction of the flux may be shifting out of the optical bands into the infrared. Similar behavior has been discussed in the case of SN 2011fe (Kerzendorf et al. 2014) and SN 2003hv (Leloudas et al. 2009). After t ∼600 days, a slower decay can be identified in all the $F475W$, $F606W$, and $F775W$-bandpasses.

Some observations of nearby Type Ia SNe show that their bolometric light curves at late phases follow the ^{56}Co decay channel (Cappellaro et al. 1997; Sollerman et al. 2004; Lair et al. 2006; Stritzinger & Sollerman 2007; Leloudas et al. 2009). These observations suggest that a turbulent, confining magnetic field traps the positrons, resulting in local

energy deposition (see Chan & Lingenfelter 1993; Milne et al. 1999, 2001; Penney & Hoeflich 2014). In contrast, ^{56}Co positron escape has been suggested in some cases (Milne et al. 1999, 2001). As the ejecta expand over time, the pre-configured magnetic field weakens to the point that the Larmor radius exceeds the size of the turbulence (see Penney & Hoeflich 2014).

We fit both the $F606W$-band and a 'pseudo-bolometric' light curve using Bateman's equation for the luminosity contribution of the ^{56}Co, ^{57}Co, and ^{55}Fe decay channels. The best fit to the pseudo-bolometric light curve and the $F606W$-band light curve give a mass ratio $M(^{57}\mathrm{Co})/M(^{56}\mathrm{Co}) = 0.094^{+0.017}_{-0.016}$ and $0.076^{+0.009}_{-0.008}$, respectively. Assuming the same mass ratio yields for isotopes of the same iron-group elements (see Graur et al. 2016, based on Truran et al. 1967 and Woosley et al. 1973), our measurements correspond to ∼3.5 times the $M(^{57}\mathrm{Fe})/M(^{56}\mathrm{Fe})$ ratio of the Sun (i.e., ∼0.0217, see Table 3 of Asplund et al. 2009). This is higher than the solar ratio ∼1.8 predicted for the W7 model (calculated from Table 3 of Iwamoto et al. 1999), and the solar ratio ∼1.7 predicted for the near-Chandrasekhar-mass three-dimensional delayed-detonation model N1600 (calculated from Table 2 of Seitenzahl et al. 2013). The $M(^{57}\mathrm{Fe})/M(^{56}\mathrm{Fe})$ ratio in our measurements is also higher compare to the ratios ∼2 and ∼1.1 suggested by the late-time quasi-bolometric light curve analysis on SN 2012cg (Graur et al. 2016) and SN 2011fe (Shappee et al. 2016). A higher metallicity progenitor could decrease the production of ^{56}Ni and result in a higher $M(^{57}\mathrm{Ni})/M(^{56}\mathrm{Ni})$ ratio (Seitenzahl et al. 2013). An enhancement of neutron excess due to electron captures in the deflagration wave could lead to the same effect.

It has been suggested that beyond ∼500 days in the ejecta, energy is shifted from the optical and near-infrared to the mid- and far-infrared (referred as the infrared catastrophe, Axelrod 1980, and see Fransson et al. 1996; Fransson & Jerkstrand 2015). The V or optical luminosity may not represent the actual behavior of the bolometric light curves. This has

never been observed so far in any Type Ia SNe (e.g., Sollerman et al. 2004; Leloudas et al. 2009; McCully et al. 2014; Kerzendorf et al. 2014; Graur et al. 2016; Shappee et al. 2016). However, Dimitriadis et al. (2017) suggested that the evolution of SN 2011fe, around 550 to 650 days, is consistent with both a model that allows for position/electron escape and a model allowing for a redistribution of flux from optical to the mid-far infrared. In our study, we fitted the $F606W$-band and optical bolometric luminosity after $\sim$650 days and do not consider the infrared catastrophe. Future studies based on a larger sample will be able to help distinguish these two possible scenarios.

In summary, our multi-band photometry of SN 2014J out to 985 days past the $B-$band maximum light clearly detected the flattening due to extra luminosity contributions other than the decay of ^{56}Co. We conclude that the high $M(^{57}\mathrm{Ni})/M(^{56}\mathrm{Ni})$ ratio estimated from the late-time luminosity evolution of SN 2014J favors a near-Chandrasekhar mass explosion model such as W7 of Iwamoto et al. (1999). Any significant circumstellar light echoes beyond 0.3 pc on the plane of the sky can be excluded by our astrometric analysis. The observations strongly suggest additional heating from internal conversion and Auger electrons of $^{57}\mathrm{Co}\rightarrow{}^{57}\mathrm{Fe}$; however, one should be cautious on the high mass ratio of ^{57}Ni to ^{56}Ni. Systematical uncertainties from the SED construction procedure, especially the missing information from NIR observations and the interpolation of the SED based on limited bandpass coverage should not be ignored (i.e., see Brown et al. 2016). Additionally, the reliability of approximating the bolometric luminosity evolution after $t\sim$500 days with the $F606W$-band emission requires more careful justification. Time for one more visit at $\sim$1180 days past the $B-$band maximum has already been allocated. This will provide another chance to better characterize the luminosity evolution of SN 2014J.

The authors are grateful to Dave Borncamp and the *HST* ACS team in fixing the distortion correction issues in ACS/WFC polarized images. Some of the data used in this

study were obtained from the Mikulski Archive for Space Telescopes (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-HST data is provided by the NASA Office of Space Science via grant NNX09AF08G and by other grants and contracts. This work also made use of the Weizmann interactive supernova data repository (WISeREP). The supernova research by Y. Yang, P. J. Brown, and L. Wang is supported by NSF grant AST-0708873. P. J. Brown was partially supported by a Mitchell Postdoctoral Fellowship. Y. Yang and M. Cracraft also acknowledge support from NASA/STScI through grant HST-GO-13717.001-A, grant HST-GO-13717.001-A, HST-GO-14139.001-A, and HST-GO-14663.001-A. The research of J. Maund is supported through a Royal Society University Research Fellowship. L. Wang is supported by the Strategic Priority Research Program "The Emergence of Cosmological Structures" of the Chinese Academy of Sciences, Grant No. XDB09000000. L. Wang and X. Wang are supported by the Major State Basic Research Development Program (2013CB834903), and X. Wang is also supported by the National Natural Science Foundation of China (NSFC grants 11178003 and 11325313).

Table 1. Log of Observations of SN 2014J with *HST* ACS/WFC POLV

Filter	Polarizer	Date (UT)	Exp (s)	Phase[a] (Days)	Date (UT)	Exp (s)	Phase[a] (Days)	Date (UT)	Exp (s)	Phase[a] (Days)
F475W	POL0V	2014-11-06	3×130	276.5	2015-03-25	3×400	415.6	2015-11-12	4×1040	648.5
F475W	POL120V	2014-11-06	3×130	276.5	2015-03-25	3×400	415.6	2015-11-12	4×1040	648.7
F475W	POL60V	2014-11-06	3×130	276.5	2015-03-25	3×400	415.7	2015-11-12	4×1040	648.8
F606W	POL0V	2014-11-06	2×40	276.6	2015-03-27	3×60	417.9	2015-11-12	4×311	649.0
F606W	POL120V	2014-11-06	2×40	276.6	2015-03-27	3×60	418.0	2015-11-13	4×311	649.0
F606W	POL60V	2014-11-06	2×40	276.6	2015-03-27	3×60	418.0	2015-11-13	4×311	649.1
F775W	POL0V	2014-11-06	2×30	276.6	2015-03-27	3×20	418.0	2015-11-12	4×100	648.5
F775W	POL120V	2014-11-06	1×55	276.6	2015-03-27	3×20	418.0	2015-11-12	4×100	648.7
F775W	POL60V	2014-11-06	1×55	276.6	2015-03-27	3×20	418.0	2015-11-12	4×100	648.9
F475W	POL0V	2016-04-08	4×1040	796.2	2016-10-12	4×1040	983.1			
F475W	POL120V	2016-04-08	4×1040	796.4	2016-10-12	4×1040	983.3			
F475W	POL60V	2016-04-08	4×1040	796.6	2016-10-12	4×1040	983.4			
F606W	POL0V	2016-04-08	4×311	796.8	2016-10-14	3×360	985.1			
F606W	POL120V	2016-04-08	4×311	796.8	2016-10-14	3×360	985.1			
F606W	POL60V	2016-04-08	4×311	796.9	2016-10-14	3×360	985.1			
F775W	POL0V	2016-04-08	4×100	796.2	2016-10-12	4×202	983.1			
F775W	POL120V	2016-04-08	4×100	796.4	2016-10-12	4×202	983.3			
F775W	POL60V	2016-04-08	4×100	796.6	2016-10-12	4×202	983.4			

[a]Days since B maximum on 2014 Feb. 2.0 (JD 245 6690.5).

Table 2: *HST* ACS/WFC late-time Photometry of SN 2014J

Filter	$F475W$		$F606W$		$F775W$	
Visit	Phase[a]	AB Magnitude	Phase[a]	AB Magnitude	Phase[a]	AB Magnitude
1	276.5	17.363±0.002	276.6	17.429±0.002	276.6	16.742±0.002
2	415.6	19.464±0.002	418.0	19.602±0.004	418.0	18.276±0.005
3	648.7	22.363±0.006	649.0	21.962±0.008	648.7	21.427±0.009
4	796.4	23.266±0.011	796.8	22.917±0.017	796.4	22.492±0.019
5	983.3	24.169±0.027	985.1	23.936±0.047	983.3	23.294±0.027

Table 3. *HST* Late-time light curve decline rate of SN 2014J

Period[a]\Filter (Days)	$F475W$ (Δmag/100 days)	$F606W$ (Δmag/100 days)	$F775W$ (Δmag/100 days)
277 – 416	1.510±0.002	1.537±0.003	1.085±0.003
416 – 649	1.244±0.003	1.021±0.004	1.366±0.004
649 – 796	0.611±0.009	0.646±0.013	0.721±0.014
796 – 983	0.483±0.016	0.540±0.026	0.429±0.018

[a]Approximate days after B maximum, 2014 Feb. 2.0 (JD 245 6690.5).



Fig. 1.— *HST* ACS/WFC $F606W$ (upper panels) and associated $F606W - F555W$ (lower panels) images of SN 2014J obtained in different visits as labeled. Each square measures $3.''2 = 54$ pc along its sides (north is up, east is left). The distance between little tick marks corresponds to $0.''1$.



Fig. 2.— The constructed late-time SED for SN 2014J. Dots show the bandpass monochromatic flux from *HST* observations at their effective wavelengths. Solid lines show the spectra with the warping procedure (left panel) and with the gray scaling (right panel) as described in Section 3. The lower panels present the total bandpass throughput curve (*HST* + ACS) for our $F475W$, $F606W$, and $F775W$ observations, showing the spectral response corresponding to the monochromatic fluxes calculated from the observed photometry.



Fig. 3.— Luminousity evolution of the monochromatic fluxes from the broadband observations (left panel) and pseudo-bolometric flux (right panel) with possible mechanisms explaining the flattening of the light curves of SN 2014J. The left panel presents the fitting and residuals of V3, V4, and V5 based on $F606W$-band observations while the right panel shows a similar plot based on the constructed pseudo-bolometric luminosity. The $F475W$ and $F775W$-band observations in the left panel and the $F606W$-band observations in the right panel are shown for comparison. The $F606W$-band observations after ∼500 days have been assumed to be proportional to the bolometric light curves (Milne et al. 2001) and free from possible γ-ray photons. Only observations after 500 days have been fitted with models accounting for all the listed isotopes or ^{56}Co plus a faint, unresolved light echo. One more visit around day 1180 will be obtained: the scheduled date of observations is marked by vertical dotted lines.



Fig. 4.— Astrometric comparisons of different visits. The x and y-axis are the position of bright sources relative to SN 2014J, most of which are HII regions in M82. The gray and black arrows are the relative motion between different visits prior to and after a linear regression with the RA and Dec. A 1-σ displacement calculated based on all the presented sources and scales are provided at the bottom right of each panel. No significant positional drift of the SN is found among all the cases, suggesting the absence of any circumstellar light echoes around 1 light year from the SN on the plane of the sky.

REFERENCES

Amanullah, R., Goobar, A., Johansson, J., et al. 2014, ApJL, 788, L21

Arnett, W. D. 1979, ApJL, 230, L37

—. 1982, ApJ, 253, 785

Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, ARAA, 47, 481

Axelrod, T. S. 1980, PhD thesis, California Univ., Santa Cruz.

Biretta, J., Kozhurina-Platais, V., Boffi, F., Sparks, W., & Walsh, J. 2004, ACS Polarization Calibration - I. Introduction and Status Report, Tech. rep.

Bonanos, A. Z., & Boumis, P. 2016, A&A, 585, A19

Brown, P. J., Breeveld, A., Roming, P. W. A., & Siegel, M. 2016, AJ, 152, 102

Brown, P. J., Smitka, M. T., Wang, L., et al. 2015, ApJ, 805, 74

Cappellaro, E., Mazzali, P. A., Benetti, S., et al. 1997, A&A, 328, 203

Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, ApJ, 345, 245

Chan, K.-W., & Lingenfelter, R. E. 1993, ApJ, 405, 614

Churazov, E., Sunyaev, R., Isern, J., et al. 2014, Nature, 512, 406

Cracraft, M., & Sparks, W. B. 2007, ACS Polarization Calibration - Data, Throughput, and Multidrizzle Weighting Schemes, Instrument Science Report ACS 2007-10, Tech. rep.

Dalcanton, J. J., Williams, B. F., Seth, A. C., et al. 2009, ApJS, 183, 67

Denisenko, D., Gorbovskoy, E., Lipunov, V., et al. 2014, The Astronomer's Telegram, 5795

Diehl, R., Siegert, T., Hillebrandt, W., et al. 2015, A&A, 574, A72

Dimitriadis, G., Sullivan, M., Kerzendorf, W., et al. 2017, ArXiv e-prints, arXiv:1701.07267

Foley, R. J., Fox, O. D., McCully, C., et al. 2014, MNRAS, 443, 2887

Fossey, J., Cooke, B., Pollack, G., Wilde, M., & Wright, T. 2014, Central Bureau Electronic Telegrams, 3792, 1

Fransson, C., Houck, J., & Kozma, C. 1996, in IAU Colloq. 145: Supernovae and Supernova Remnants, ed. T. S. Kuhn, 211

Fransson, C., & Jerkstrand, A. 2015, ApJL, 814, L2

Gao, J., Jiang, B. W., Li, A., Li, J., & Wang, X. 2015, ApJL, 807, L26

Gerke, J. R., Kochanek, C. S., & Stanek, K. Z. 2014, The Astronomer's Telegram, 5808

Gonzaga, S., Hack, W., Fruchter, A., & Mack, J. 2012, The DrizzlePac Handbook (Baltimore: STScI)

Goobar, A., Johansson, J., Amanullah, R., et al. 2014, ApJL, 784, L12

Graham, M. L., Nugent, P. E., Sullivan, M., et al. 2015, MNRAS, 454, 1948

Graur, O., Zurek, D., Shara, M. M., et al. 2016, ApJ, 819, 31

Hartig, G. F. 2009, WFC3 SMOV Programs 11436/8: UVIS On-orbit PSF Evaluation, Instrument Science Report WFC3 2009-38, Tech. rep., Baltimore: STScI

Hoyle, F., & Fowler, W. A. 1960, ApJ, 132, 565

Itagaki, K., Kaneda, H., Yamaoka, H., et al. 2014, Central Bureau Electronic Telegrams, 3792

Iwamoto, K., Brachwitz, F., Nomoto, K., et al. 1999, ApJS, 125, 439

Johansson, J., Goobar, A., Kasliwal, M. M., et al. 2014, ArXiv e-prints, arXiv:1411.3332

Kerzendorf, W. E., Taubenberger, S., Seitenzahl, I. R., & Ruiter, A. J. 2014, ApJL, 796, L26

Khokhlov, A. M. 1991, A&A, 245, 114

Lair, J. C., Leising, M. D., Milne, P. A., & Williams, G. G. 2006, AJ, 132, 2024

Leloudas, G., Stritzinger, M. D., Sollerman, J., et al. 2009, A&A, 505, 265

Lundqvist, P., Nyholm, A., Taddia, F., et al. 2015, A&A, 577, A39

Ma, B., Wei, P., Shang, Z., Wang, L., & Wang, X. 2014, The Astronomer's Telegram, 5794

McCully, C., Jha, S. W., Foley, R. J., et al. 2014, ApJ, 786, 134

Milne, P. A., The, L.-S., & Leising, M. D. 1999, ApJS, 124, 503

—. 2001, ApJ, 559, 1019

Ohlmann, S. T., Kromer, M., Fink, M., et al. 2014, A&A, 572, A57

Pakmor, R., Hachinger, S., Röpke, F. K., & Hillebrandt, W. 2011, A&A, 528, A117

Pakmor, R., Kromer, M., Taubenberger, S., et al. 2012, ApJL, 747, L10

Penney, R., & Hoeflich, P. 2014, ApJ, 795, 84

Perlmutter, S., Aldering, G., Goldhaber, G., et al. 1999, ApJ, 517, 565

Riess, A. G., Filippenko, A. V., Challis, P., et al. 1998, AJ, 116, 1009

Riess, A. G., Macri, L. M., Hoffmann, S. L., et al. 2016, ApJ, 826, 56

Röpke, F. K., Kromer, M., Seitenzahl, I. R., et al. 2012, ApJL, 750, L19

Schlafly, E. F., & Finkbeiner, D. P. 2011, ApJ, 737, 103

Seitenzahl, I. R., Taubenberger, S., & Sim, S. A. 2009, MNRaS, 400, 531

Seitenzahl, I. R., Timmes, F. X., & Magkotsios, G. 2014, ApJ, 792, 10

Seitenzahl, I. R., Ciaraldi-Schoolmann, F., Röpke, F. K., et al. 2013, MNRAS, 429, 1156

Shappee, B. J., Stanek, K. Z., Kochanek, C. S., & Garnavich, P. M. 2016, ArXiv e-prints, arXiv:1608.01155

Sirianni, M., Jee, M. J., Benítez, N., et al. 2005, PASP, 117, 1049

Sollerman, J., Lindahl, J., Kozma, C., et al. 2004, A&A, 428, 555

Sparks, W. B., & Axon, D. J. 1999, PASP, 111, 1298

Srivastav, S., Ninan, J. P., Kumar, B., et al. 2016, MNRAS, 457, 1000

Stritzinger, M., & Sollerman, J. 2007, A&A, 470, L1

Swartz, D. A., & Wheeler, J. C. 1991, ApJL, 379, L13

Taubenberger, S., Elias-Rosa, N., Kerzendorf, W. E., et al. 2015, MNRAS, 448, L48

Truran, J. W., Arnett, W. D., & Cameron, A. G. W. 1967, Canadian Journal of Physics, 45, 2315

Woosley, S. E., Arnett, W. D., & Clayton, D. D. 1973, ApJS, 26, 231

Yang, Y., Wang, L., Baade, D., et al. 2017, ApJ, 834, 60

Zheng, W., Shivvers, I., Filippenko, A. V., et al. 2014, ApJL, 783, L24

This manuscript was prepared with the AAS LaTeX macros v5.2.